





(a) $\uparrow E_k$



(b) $\uparrow E_k$







energy spectrum. Such a behavior is dictated by the selection rules and energy conservation law: Direct optical transitions in QWs of critical thickness are allowed only between branches 1 and 2 and 3 and 4 and these transitions can occur at wave vector, which leads to the emergence of extremum at $k = 2\pi/L$. In QWs of noncritical thickness [Fig. 5(b)], the branches 2 and 3 anticross at a finite wave vector, and direct optical transitions between them become allowed. It leads to the emergence of an additional sharp band in the absorption spectrum at $k = 2\pi/L$. The spectral shape of this absorption band is determined by the van Hove singularities in the density of states.

To summarize, we have described the splitting of Dirac states in HgTe/CdTe quantum wells of critical and close-to-critical thicknesses. In structures of critical thickness, the splitting between the Dirac cones reaches a value of 15 meV and is dominated by symmetry-enforced light-hole and heavy-hole mixing at the quantum well interfaces. These structures

Financial support by the RFBR, RF President Grants No. MD-3098.2014.2 and No. NSh-1085.2014.2, EU project

SPANGL4Q, and the Ó Dynasty Ó Foundation is gratefully acknowledged. Work at the University of Colorado, Boulder by A.Z. was supported by the US Department of Energy, Office of Science, Basic Energy Science, Materials Sciences and Engineering Division under Contract No. DE-FG02-93ER46959 to the University of Colorado.

[1] M. Z. Hasan and C. L. Kane, *Rev. Mod. Phys.* **82**, 3045 (2010).

[2] X.-L. Qi and S. C. Zhang, *Rev. Mod. Phys.* **83**, 1057 (2011).

[3] Y. Xia, D. Qian, D. Hsieh, L. Wray, A. Pal, H. Lin, A. Bansil, D. Grauer, Y. AP, R. G. 6(.-)2440 I oJ.6(.-)24406(C)-.5(6)20.3(v)1521(and, 2440 I od)-25028(a)M.5(.-)24806(C)Z.4(s)